



Effects of single-session versus multi-session non-invasive brain stimulation on craving and consumption in individuals with drug addiction, eating disorders or obesity: A meta-analysis



Sensen Song^{a, b, 1}, Anna Zilverstand^{c, 1}, Wenjun Gui^{d, f}, Hui-jie Li^{e, f}, Xiaolin Zhou^{a, b, g, h, *}

^a School of Humanities, Tongji University, Shanghai, 200092, China

^b School of Psychological and Cognitive Sciences, Peking University, Beijing, 100871, China

^c Department of Psychiatry, University of Minnesota, Minneapolis, MN, 55414, USA

^d Key Laboratory of Mental Health, Institute of Psychology, Chinese Academy of Sciences, Beijing, 100101, China

^e CAS Key Laboratory of Behavioral Science, Institute of Psychology, Chinese Academy of Sciences, Beijing, 100101, China

^f Department of Psychology, University of Chinese Academy of Sciences, Beijing, 100101, China

^g Beijing Key Laboratory of Behavior and Mental Health, Peking University, Beijing, 100871, China

^h PKU-IDG/McGovern Institute for Brain Research, Peking University, Beijing, 100871, China

ARTICLE INFO

Article history:

Received 15 June 2018

Received in revised form

18 December 2018

Accepted 24 December 2018

Available online 27 December 2018

Keywords:

Brain stimulation

tDCS

rTMS

dIPFC

Addiction

Eating disorders

ABSTRACT

Background: Brain stimulation interventions are increasingly used to reduce craving and consumption in individuals with drug addiction or excessive eating behavior. However, the efficacy of these novel treatments and whether effect sizes are affected by the length of the intervention has not been comprehensively evaluated.

Objective: A meta-analytical approach was employed to evaluate the effectiveness of non-invasive excitatory brain stimulation [transcranial Direct Current Stimulation (tDCS) and high-frequency repetitive Transcranial Magnetic Stimulation (rTMS)] targeted at dorsolateral prefrontal cortex (dlPFC) for reducing craving and consumption levels in drug and eating addiction, including both single- and multi-session protocols.

Methods: After a comprehensive literature search, 48 peer-reviewed studies (1095 participants in total) were included in the current meta-analysis. We computed Hedge's *g* as a conservative measure for evaluating effect sizes.

Results: Random effects analyses revealed a small effect of neuromodulation interventions on craving and a medium effect on consumption, favoring active over sham stimulation. These effects did not differ across the different populations investigated (alcohol, nicotine, illicit drugs, eating addictions) or by the used technique (rTMS/tDCS, left/right hemisphere). Multi-session protocols showed a larger effect size for reducing craving and consumption than single-session protocols, with a positive linear association between the number of sessions or administered pulses and craving reduction, indicating a dose-response effect.

Conclusions: Our results provide compelling evidence that novel non-invasive brain stimulation targeted at dlPFC reduces craving and consumption levels (providing the first meta-analytical evidence for the latter effect in drug addiction), with larger effects in multi-session as compared to single-session interventions.

© 2018 Elsevier Inc. All rights reserved.

Introduction

Substance use disorders, eating disorders and obesity are major public health problems, which are costly and pervasive. According to 2016 National Survey on Drug Use and Health [1], more than 20.1 million (7.4%) adolescents or adults in the United States experienced substance use disorder, while 37.9% of adults aged 20 years

* Corresponding author. School of Psychological and Cognitive Sciences, Peking University, 5 Yiheyuan Road, Beijing, 100871, China.

E-mail address: xz104@pku.edu.cn (X. Zhou).

¹ Sensen Song and Anna Zilverstand contributed equally to this article.

and over were defined as obese (Health United States Report 2016; [2]), with a cumulative lifetime risk of eating disorders characterized by excessive eating of 5.7% of the population (National Comorbidity Survey Replication 2001–2003; [3]). Novel treatment approaches, such as neuromodulation approaches, are urgently needed, as the efficacy of currently available therapeutic interventions in preventing relapse in drug addiction is only 40–60% [4], with similarly low success rates (~50%) in eating disorders [5].

The hallmark of drug addiction and excessive eating behavior is the lack of ability to refrain from consumption (e.g. smoking, drinking or eating); even when consumption is associated with potentially negative consequences. Compulsive consumption is generally preceded by craving, which has been defined as an “intensive desire or urge” to consume in drug addiction [6]. In drug addiction, craving predicts the maintenance of drug seeking behavior, as well as relapse after abstinence [7]. Reducing craving has hence been proposed to be a meaningful goal for clinical intervention studies [7]. Similar to drug craving, food craving is defined as an intense desire to consume a particular food that is difficult to resist [8]. Although still controversial, the existing evidence similarly points to food craving as a critical factor in the maintenance of eating disorders and obesity [9], with craving predicting body mass index and consumption of high caloric foods such as sweets and high-fat food [10]. Both drug addiction and disorders of excessive food intake are further characterized by a break-down of inhibitory self-control, which aggravates the intense urge to consume [11–13]. Because of these observed parallels between drug addiction and excessive eating behavior, the later has increasingly been conceptualized as eating addiction [14–16], which encompasses pathological binge eating as well as uncontrolled eating in individuals without a clinical diagnosis [16]. This comparison has been further substantiated by findings implicating common neural mechanisms between eating and drug addictions [12,13,16].

The intensified urge to consume in drug addiction is thought to have its neurobiological substrate both in an increased phasic (yet reduced tonic) response of the dopaminergic reward system in the presence of drugs or drug cues [17–21] and the decreased engagement of prefrontal systems underlying inhibitory self-control, such as the dorsolateral prefrontal cortex (dlPFC) [18,22]. Studies across different addicted populations have linked increased activation levels in the dopaminergic reward system during drug cue exposure to increased subjective craving [23,24]. In parallel, in eating addiction, the sensitization of the dopaminergic reward system has been linked to an increased drive for excessive intake of food [12,13,16]. Aggravating the problem, this observed hyper-reactivity of the reward system is potentiated by an reduced engagement of prefrontal systems (including the dlPFC) during attempts to self-regulate craving and inhibit consumption of drugs [22,25,26] or food [12,13,16]. The primary role of the dlPFC is thought to be the representation of contextual information when faced with conflicting outcomes [27], hence fulfilling a crucial role in the selection (and inhibition) of behavioral and cognitive responses in the face of different options [28–30]. Evidence further suggests that the dlPFC is the apex of the frontal control hierarchy, exerting influence on other frontal brain regions through its broad efferent connections [30]. As such its recruitment may be necessary for successful inhibition of a behavioral (consumption) or cognitive (craving) response. In drug addiction, the dlPFC has been shown to be hypo-active, specifically when attempting to inhibit behavioral responses [26,31] or to cognitively self-regulate craving [26,32,33], while it is hyperactivated when not attempting to self-regulate craving during drug cue exposure [26]. In analogy to its role in drug addiction, the dlPFC has also been implicated in the failure to inhibit excessive food intake in eating disorders and obesity [12,13].

Moreover, the attenuation of dlPFC activity by inhibitory forms of rTMS led to overconsumption of high-caloric foods in healthy weight individuals, providing evidence for its causal role in reducing consumption [34,35].

Recent reviews of the neurobiological effects of the available cognitive-behavioral and pharmacological treatments in drug addiction have proposed that the dlPFC is an important neural target of treatment, showing the potential for a (partial) normalization of its functions after treatment [26,32]. While the dlPFC is by no means the only target for treatment and inhibitory control is not the only mechanism impaired in drug and eating addictions (for in depth reviews see Refs. [12,13,16–18,20,26]), the dlPFC is particularly suited as a target for non-invasive neuromodulation approaches with currently widely used non-invasive neuromodulation techniques, such as repetitive Transcranial Magnetic Stimulation (rTMS) and transcranial Direct Current Stimulation (tDCS), both because of its involvement in inhibitory control and its closeness to the scalp [36]. Repetitive TMS with standard figure-eight electromagnetic coils exerts its neuromodulatory influence by generating (repetitive) magnetic pulses which induce small electrical currents within a focal area in the superficial brain tissue below the scalp [36]. In rTMS, trains of pulses are used to achieve a sustained alteration of neural excitability in the targeted region [37]. Excitatory rTMS, which has a frequency of more than 5 Hz (“high frequency rTMS”) lowers the threshold for (self-) engaging the targeted brain region, whereas low frequency stimulation (≤ 1 Hz) would have inhibitory effects [38]. Similarly, tDCS can either increase or decrease neuronal excitability. tDCS generates a low intensity electric field (0.5–2 mA) between two electrodes, increasing neuronal excitability under the anodal, but decreasing it under the cathodal electrode [39], allowing for a targeted approach. When applied for a sufficient duration, increases in cortical excitability are sustained [39] and linked to changes in synaptic plasticity [40]. The hypothesized neuro-behavioral mechanism of the reviewed excitatory rTMS/tDCS interventions is thus to upregulate neuronal excitability of the dlPFC, such that the threshold for engaging this region during (self-)regulation of craving and consumption is lowered. In this meta-analysis, we focused on quantifying the effects of excitatory neuromodulation targeted at dlPFC, because we hypothesized that such interventions are of particular clinical benefit in a population characterized by hypoactivation of the dlPFC during attempted self-regulation.

There are a number of reasons for us to perform the current meta-analysis. First, no meta-analysis has been carried out to quantify the effects of neuromodulation approaches to reduce both craving and consumption in populations with drug addiction and excessive eating behavior. Previous meta-analyses on this topic only focused on craving measures of these disorders [41], or only concerned the regulation of eating behavior [42,43] or two types of drug users (alcohol and nicotine) [44]. These previous analyses included a much smaller number of studies (17 studies [41], 11 studies [42], or 10 studies [44], as compared to 48 studies in the current analysis), providing a less reliable estimate of the effect sizes. Second, none of the previous meta-analyses investigated the impact of the number of sessions or administered pulses, which could be important when evaluating a potential dose-response effect. However, previous findings from individual studies that directly compared single-to multiple-session protocols found that single-session interventions yielded smaller improvements as compared to interventions with five [45,46] or eight sessions [47]. Third, previous meta-analyses on drug addiction only used craving as an outcome measure [41,44], even though reducing consumption is the most important goal of these interventions. Fourth, previous meta-analyses on the effects in drug addiction only

included individuals with alcohol and nicotine addictions [41,44], but not illicit drug users. Fifth, the current meta-analysis addressed potential influences by differences in methodological approach (stimulated hemisphere, study design) on both craving and consumption, in addition to the re-evaluation of the effect of the stimulation technique employed (rTMS vs tDCS) [41].

Methods

To identify pertinent studies, a two-staged literature search was carried out. First, an online search was conducted in PubMed, Web of Science, PsycINFO and Embase database covering peer-reviewed articles from January 1990 to July 2018. This search was performed following the PICO-method (Patient/Population, Intervention, Comparison, Outcome). Key elements were the 'P' (substance or eating addiction), the 'I' (tDCS or rTMS stimulation), the 'C' (Active and Sham stimulation) and the 'O' (craving or consumption levels). See *Supplementary A* for full search terms. Second, an additional literature search was performed using the reference lists of the identified studies, of other meta-analytical studies [41–44] and a number of relevant review articles [36,48–53]; the goal was to search for as many potential studies as possible.

Inclusion criteria

This work followed PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) guidelines. Peer-reviewed studies published in all language journals and meeting the following criteria were included in the current meta-analysis: (1) randomized controlled trials (RCTs) and controlled clinical trials (CCTs), focusing on high-frequency rTMS or anodal tDCS targeting the dlPFC in patients with (a) substance use disorder (e.g. alcohol, cannabis, cocaine, heroin, methamphetamine and nicotine) or frequent smoking or (b) eating disorders (Bulimia Nervosa/Binge Eating type) or individuals with frequent food craving or obesity; (2) including at least one craving or consumption measure as an outcome; (3) comparing real to sham brain stimulation; (4) having provided means, standard deviations (SDs), *t*, *F* or *p* statistics and the number of participants in each intervention group, or other data that could be used to calculate effect size. The inclusion criteria did not consider the craving assessment tool, consumption assessment tool, number of stimulation sessions, stimulated hemisphere, or method of localization for the site of stimulation.

Exclusion criteria

We excluded studies from our meta-analysis if they: (1) focused on populations other than the ones mentioned above (e.g. schizophrenia, Prader-Willi syndrome, depression, posttraumatic stress disorder or chronic pain); (2) assessed the effects of stimulation using techniques other than tDCS or high frequency rTMS (e.g., deep TMS, continuous theta burst stimulation, electro convulsive therapy and low frequency rTMS); (3) assessed the effect of dlPFC stimulation using outcome measures other than craving or consumption; (4) lacked a sham stimulation condition in experiment design; (5) did not provide enough information to calculate effect size; (6) were case studies or review articles.

Data extraction

The extracted data included the study name, population, type of substance involved, number of stimulation sessions per condition, side of stimulation (right vs. left), stimulation technique (tDCS vs. rTMS), number of participants, study design, blinding method, blinding evaluation, total number of pulses, craving measures,

consumption measures and standardized effect sizes for the effect of stimulation on craving and consumption levels. When means and standard deviations were not available, effect sizes were computed from *t*-values, *F*-values or *p*-values. When studies included more than one measures on craving or consumption, we averaged them to compute one pooled effect size. For example, if a study included two measures of craving [(Visual Analogue Scale (VAS) and Food Craving Questionnaire State (FCQ-S)], these two measurements were summarized by their pooled effect size [54]. If a study reported results of measures both during the stimulation and post-stimulation, only the data of the post-stimulation was included [55]. For the five studies [56–60], which included both left and right dlPFC stimulation data, a combined effect size for both sites was calculated. However, when evaluating left with right hemisphere stimulation, we computed separate effect sizes for each stimulation site for these studies. When the reported data was insufficient for the current data analysis, the authors were contacted. If the authors could not be reached or the data were only available in chart format, means and SDs were estimated using Engauge Digitizer [61]. Each study included in the meta-analysis was independently coded by two of the authors (S.S. and W.G.); discrepancies were addressed by discussion.

Data analysis

To assess the risk of bias within the individual studies, we used a standardized critical appraisal instrument named the Cochrane Collaboration's risk of bias tool [62]. Ratings (high, low, or unclear risk) were assigned to evaluate bias on the following domains: random sequence generation (selection bias), allocation concealment (selection bias), blinding of participants and personnel (performance bias), blinding of outcome assessment (detection bias), incomplete outcome data (attrition bias), selective reporting (reporting bias) and other bias (if any). In addition, we evaluated the used control condition and blinding procedures for all included studies. Blinding of the outcome assessment was assessed for craving and consumption independently. Two reviewers (S.S. and W.G.) independently evaluated and recorded their judgments, and any disagreements were resolved by discussion.

To examine the magnitude of the effect of non-invasive brain stimulation on craving and consumption levels, we computed a synthesized effect size Hedge's *g* with a corresponding confidence interval (CI) of 95%. Hedge's *g* is considered to be a conservative estimate, which is useful for studies with small sample sizes [63]. The magnitude of *g* is conventionally interpreted as reflecting a small (if $g = 0.2-0.5$), medium (if $g = 0.5-0.8$), or a large effect size (if $g > 0.8$) [63]. The direction of the effect size was defined as positive if an active stimulation condition was better than the sham condition on craving or consumption measures. A random-effects model was used for all between-group analyses, which is in accordance with the underlying assumption of the non-homogeneity between individual study samples. Compared to a fixed-effects model, the random-effects model provides a more conservative estimate of precision and is more appropriate for generalization beyond the included studies [63,64]. We first calculated the overall effect size (independent of the number of sessions) on craving and consumption levels (for the 5 studies reporting both single-session and multi-session effects [45–47,65,66], only the multi-session effect was included here), and then estimated the effect sizes of single-session and multi-session interventions separately. To compare single-session with multi-session effects, a *Q* test was conducted [67]. We further investigated the influence of the number of sessions (or the total number of pulses) on both craving and consumption using a moderator variable analysis [67]. In addition, we assessed the

potential influences of the population, stimulation technique and study design on such intervention effects. In particular, as five studies [56–60] included both left and right dlPFC stimulation data, in order to exclude the multiple effect sizes in individual studies, we used Z-test (instead of Q-test) to assess the potential influences of the stimulated hemisphere [63]. Furthermore, we used Egger's regression intercept test to measure for publication bias [68]. All suitable data were calculated with the software Comprehensive Meta-Analysis 2.0 (CMA) (<http://www.meta-analysis 2.0.com>).

Results

The results of the initial search are summarized in Fig. 1. We included a total of 48 studies focusing on non-invasive brain stimulation interventions in drug or eating addiction. The characteristics of these studies are summarized in Table 1. Not all studies reported both craving and consumption measures, and only 5 studies evaluated both a single-session and a multi-session effect (Supplementary Table 1). Specifically, we included 44 articles on studies assessing the overall intervention effect of brain stimulation on craving (33 with single-session and 15 with multi-session interventions). In addition, we included 15 articles on studies that investigated the overall intervention effect of brain stimulation on consumption (10 with single-session and 7 with multi-session interventions). All included studies used excitatory rTMS or tDCS stimulation.

Assessment of risk of bias in the included studies

The evaluation of the risk of bias for each included study was summarized in Supplementary Table 2. The results indicated that overall the included studies were of high quality (i.e., at relatively low risk of bias). In addition, the evaluation of the control condition and blinding procedures for all included studies was summarized in Supplementary Table 3. Both evaluations found that all included studies used appropriate procedures to avoid bias. Only three studies [69–71] reported a partial failure of the employed blinding procedures (e.g., Goldman et al., 2011, in which the authors

reported that “participants were able to identify whether they were receiving real or sham tDCS at a rate better than chance.”). To ensure that these three studies did not bias our findings, we repeated all analyses while excluding these studies (see below).

Craving: overall intervention effect and the influence of multiple sessions and session length

We found a significant overall effect (independent of the number of sessions) of non-invasive neuromodulation on craving levels with a small effect size ($g = 0.456$; CI: 0.328–0.583), favoring active over sham stimulation in reducing craving ($z = 7.009$, $p < 0.0001$) (Fig. 2). The Egger's test indicated that there was no publication bias ($t_{(42)} = 0.982$, $p = 0.332$). Single-session interventions had a small pooled standardized effect ($g = 0.360$; CI: 0.225–0.495), indicating that active stimulation was more effective than sham stimulation in reducing craving ($z = 5.211$, $p < 0.0001$) (Fig. 3). More importantly, multi-session interventions had a medium pooled standardized effect size ($g = 0.683$; CI: 0.401–0.964), again indicating that active stimulation was more effective than sham stimulation ($z = 4.751$, $p < 0.0001$) (Fig. 4). Furthermore, Q-test analysis revealed that the effect size of multi-session protocols was significantly larger than the effect size of single session protocols ($Q = 4.138$, $p = 0.042$), indicating that multi-session interventions were more beneficial for the reduction of craving as compared to single-session interventions. Finally, the moderator analysis found a significant positive linear relationship between the number of sessions and craving levels ($\beta = 0.045$, CI: 0.010–0.081, $Q = 6.364$, $p = 0.012$, Supplementary Fig. 1), and a significant positive linear relationship between the total number of pulses (including the 18 rTMS studies which have this parameter) and the craving level ($\beta = 0.0001$, CI: 0.00002–0.00013, $Q = 8.465$, $p = 0.004$, Fig. 5).

To investigate if neuromodulation interventions were equally effective in the potentially more severely impaired populations with a clinical diagnosis of either an eating disorder or substance addiction, we conducted a supplementary analysis, excluding the 16 studies performed in populations without an eating disorder diagnosis (individuals with obesity or excessive food craving, Supplementary B). We obtained the same results as above, replicating the overall effect of neuromodulation, the larger effect after multiple sessions and the positive linear relationship between the number of sessions/pulses and effect size (Supplementary B). Additionally, we repeated all analyses after excluding the three studies which reported failure of blinding procedures [69–71], also replicating all the findings above (Supplementary C).

Consumption: overall intervention effect and the influence of multiple sessions and session length

We found a significant overall effect (independent of the number of sessions) of non-invasive neuromodulation on consumption levels with a medium pooled standardized effect size ($g = 0.657$; CI: 0.390–0.924), favoring active stimulation over sham stimulation ($z = 4.819$, $p < 0.0001$) (Fig. 6). Egger's test again indicated that there was no publication bias ($t_{(13)} = 0.736$, $p = 0.475$). Single-session interventions again had a small pooled standardized effect ($g = 0.433$; CI: 0.128–0.738), indicating that active stimulation was more effective than sham stimulation in reducing consumption ($z = 2.786$, $p = 0.005$) (Fig. 7), while multi-session interventions in contrast showed a large pooled standardized effect size ($g = 0.950$; CI: 0.625–1.275) for active stimulation as compared to sham stimulation ($z = 5.734$, $p < 0.0001$) (Fig. 8). Moreover, Q-test analysis revealed that multi-session protocols for reducing consumption levels had a significantly larger effect size than single session protocols ($Q = 5.176$, $p = 0.023$), again indicating that multi-session




Fig. 1. Flow chart of the study selection process. Reasons for exclusion included the use of stimulation techniques other than tDCS or high frequency rTMS (such as continuous theta burst stimulation, electro convulsive therapy and low frequency rTMS) or the investigation of effects in other clinical populations (such as Schizophrenia, Depression, Posttraumatic Stress Disorder or chronic pain).

Table 1
Study and sample characteristics for included studies.

First author	Study population	Number of sessions (per condition)	Anodal Stimulation side (dIPFC)	Stimulation Technique	Participants number	Study Design	total number of pulses (per condition)	Craving measure	Consumption measure
Alcohol (12 studies)									
Boggio et al.(2008)	Alcohol dependence	1	Both	tDCS	13	W	NA	AUQ *	NA
da Silva et al.(2013)	Alcohol dependence	5	Left	tDCS	13	B	NA	OCDS *	NA
den Uyl et al. (2015)	Alcohol use disorder	1	Left	tDCS	26	B	NA	AAAQ *	NA
Herremans et al.(2012)	Alcohol dependence	1	Right	rTMS	31	B	1560	OCDS	NA
Herremans et al.(2013)	Alcohol dependence	1	Right	rTMS	29	W	1560	OCDS	NA
Herremans et al.(2015)	Alcohol dependence	1	Right	rTMS	24	W	1560	VAS	NA
Höppner et al.(2011)	Alcohol dependence	10	Left	rTMS	19	B	NA	OCDS	NA
Klauss et al.(2014)	Alcohol dependence	5	Right	tDCS	33	B	NA	OCDS	NA
Klauss et al.(2018)	Alcohol dependence	10	Right	tDCS	45	B	NA	OCDS	NA
Mishra et al.(2010)	Alcohol dependence	10	Right	rTMS	45	B	10000	ACQ *	NA
Nakamura-Palacios et al.(2012)	Alcohol dependence	1	Left	tDCS	32	W	NA	OCDS *	NA
Wietschorke et al.(2016)	Alcohol dependence	1	Right	tDCS	30	B	NA	VAS *	NA
Food (20 studies)									
Barth et al. (2011)	Healthy women who endorsed frequent food cravings	1	Left	rTMS	10	W	3000	VAS	NA
Bravo et al.(2016)	Obesity	5	Right	tDCS	10	B	NA	VAS	NA
Burgess et al.(2016)	Binge Eating Disorder (also subthreshold)	1	Right	tDCS	30	W	NA	VAS *	Calories consumed *
Claudino et al. (2011)	Bulimia Nervosa or Eating Disorder not otherwise specified	1	Left	rTMS	22	B	1000	VAS and FCQ-S *	NA
Fregni et al. (2008a)	Healthy subjects with frequent food cravings	1	Both	tDCS	21	W	NA	VAS *	Calories consumed *
Gluck et al.(2015)	Obesity	3	Left	tDCS	9	B	NA	NA	Calories consumed
Gluck et al.(2017)	Obesity	15	Left	tDCS	20	B	NA	NA	Calories consumed *
Goldman et al. (2011)	Healthy subjects with frequent food cravings	1	Right	tDCS	19	W	NA	VAS	Food consumed
Grundeis et al.(2017)	Obesity	1	Left	tDCS	23	W	NA	NA	Calories consumed
Heinitz et al.(2017)	Obesity	15	Left	tDCS	29	B	NA	NA	Calories consumed *
Jauch-Chara et al.(2014)	Healthy men with low cognitive restraint for food consumption	Both 1 and 8	Right	tDCS	14	W	NA	VAS *	Calories consumed *
Kekic et al.(2014)	Healthy women with frequent food cravings	1	Right	tDCS	20	W	NA	FCQ-S	Food consumed
Kekic et al.(2017)	Bulimia Nervosa	1	Both	tDCS	39	W	NA	VAS *	NA
Kim et al.(2018)	Obesity	4	Left	rTMS	57	B	4000	VAS	NA
Lapenta et al. (2014)	Healthy women with frequent food cravings	1	Right	tDCS	9	W	NA	VAS *	Calories consumed *
Ljubisavljevic et al.(2016)	Healthy individuals with frequent food cravings	5	Right	tDCS	27	B	NA	FCQ-S and FCI *	NA
Montenegro et al. (2012)	Overweight	1	Left	tDCS	9	W	NA	VAS *	NA
Ray et al.(2017)	Obesity	1	Right	tDCS	18	W	NA	VAS	Calories consumed
Uher et al. (2005)	Women with strong food cravings	1	Left	rTMS	28	B	1000	VAS *	Calories consumed
Van den Eynde et al. (2010)	Bulimia Nervosa or Eating Disorder not otherwise specified	1	Left	rTMS	37	B	1000	VAS *	NA
Nicotine (9 studies)									
Amiaz et al. (2009)	Nicotine dependence	10	Left	rTMS	21	B	10000	VAS and sTCQ *	Self-report Cigarettes consumed *
Boggio et al. (2009)	Smoking ≥ 10 cigarettes per day		Left	tDCS	23	B	NA	VAS *	

		Craving: Both 1 and 5; Consumption: 5							Self-report Cigarettes consumed *
Fecteau et al.(2014)	Average daily intake of at least 15 cigarettes	Craving: 5 Consumption: both 1 and 5	Right	tDCS	12	W	NA	sQSU *	Self-report Cigarettes consumed *
Fregni et al. (2008b)	Smoking 15 or more cigarettes per day	1	Both	tDCS	24	W	NA	VAS *	NA
Johann et al. (2003)	Average daily intake of at least 18 cigarettes	1	Left	rTMS	11	W	1000	VAS*	NA
Li et al.(2013)	Nicotine dependence	1	Left	rTMS	14	W	3000	VAS *	NA
Li et al.(2017)	Nicotine dependence	1	Left	rTMS	10	W	3000	VAS	NA
Pripfl et al.(2014)	Nicotine dependence	1	Left	rTMS	11	W	1200	VAS *	NA
Xu et al. (2013)	Daily intake of at least 10 cigarettes	1	Left	tDCS	24	W	NA	UTS	NA
Drugs (7 studies)									
Batista et al.(2015)	Cocaine dependence	5	Right	tDCS	36	B	NA	OCCS *	NA
Boggio et al. (2010)	Marijuana use at least 3 times per week	1	Both	tDCS	17	B	NA	VAS *	NA
Sahlem et al.(2017)	Cannabis Use Disorder	1	Left	rTMS	16	W	4000	MCQ	NA
Shahbabaie et al. (2014)	Methamphetamine dependence	1	Right	tDCS	30	W	NA	VAS	NA
Shen et al.(2016)	Heroin dependence	Both 1 and 5	Left	rTMS	10	B	10000	VAS *	NA
Su et al.(2017)	Methamphetamine dependence	Both 1 and 5	Left	rTMS	30	B	6000	VAS *	NA
Shahbabaie et al.(2018)	Methamphetamine use disorder	1	Right	tDCS	15	W	NA	VAS *	NA

AAAQ: alcohol approach and avoidance questionnaire; ACQ: Alcohol Craving Questionnaire; AUQ: Alcohol Urge Questionnaire; B/W: between-subject design/within-subject design; dlPFC: dorsolateral prefrontal cortex; FCI: Food Craving Inventory; FCQ-S: Food Craving Questionnaire State; MCQ: Marijuana Craving Questionnaire; NA: not available; OCDS: Obsessive Compulsive Drinking Scale; rTMS: repetitive transcranial magnetic stimulation; STCQ: short version of the Tobacco Craving Questionnaire; sQSU: standardized Questionnaire of Smoking Urges; tDCS: transcranial direct current stimulation; UTS: Urge to Smoke; OCCS: Obsessive-Compulsive Cocaine Scale; VAS: Visual Analogue Scale; * represent significant craving or consumption reduction after intervention, if an article reported the effect of both single-session and multi-session, this only represent the effect of the later.




Fig. 2. The overall effect of neuromodulation on craving.

protocols were more beneficial. However, the moderator analysis testing for a positive linear association between the number of sessions and effect size ($\beta = 0.032$, CI: -0.001 – 0.008 , $Q = 2.488$, $p = 0.115$) was not significant, possibly due to the lower number of studies measuring consumption as compared to craving (Supplementary Fig. 2). The relationship between administered pulses and consumption could not be investigated since there were only two available rTMS studies. The additional analyses excluding individuals without a clinical diagnosis of eating disorder studies (Supplementary B), as well as the analyses excluding studies with partially failed blinding procedures (Supplementary C) confirmed the above findings.

Differences in effects between different study populations

The comparison of the effect sizes in different study populations (alcohol, nicotine, illicit drugs or eating addiction) revealed no significant differences in the overall effect size for the modulation of craving ($Q = 2.545$, $p = 0.467$). In addition, when investigating each population separately, we found a significant effect with small to medium effect sizes on craving level in individuals with alcohol ($g = 0.382$, CI: 0.078 – 0.685 ; $z = 2.465$, $p = 0.014$), eating ($g = 0.426$, CI: 0.261 – 0.592 ; $z = 5.047$, $p < 0.0001$), drug ($g = 0.514$, CI: 0.073 – 0.955 ; $z = 2.284$, $p = 0.022$) and nicotine ($g = 0.627$, CI: 0.407 – 0.846 ; $z = 5.601$, $p < 0.0001$) addiction. The comparison between populations of the overall effect size for a reduction of consumption level was marginally significant ($Q = 2.970$,




Fig. 3. The effect of single-session neuromodulation on craving.




Fig. 4. The effect of multi-session neuromodulation on craving.

$p = 0.085$), revealing a slightly larger effect in individuals with nicotine addiction (with consumption levels generally being measured by cigarettes smoked per day as indicated by self-report, see Table 1) as compared to eating addiction (based on consumed

food during a test after the intervention, see Table 1). When considering these two populations separately, we found a significant reduction of consumption levels in both populations, with a large effect size in smokers ($g = 1.138$, CI: 0.543–1.733; $z = 3.751$,




Fig. 5. Regression of the total number of pulses on the effect size of neuromodulation of craving. $\beta = 0.0001$, 95% CI: 0.00002–0.00013, $Q = 8.465$, $p = 0.004$.




Fig. 6. The overall effect of neuromodulation on consumption.




Fig. 7. The effect of single-session neuromodulation on consumption.

$p < 0.0001$) and a medium effect size for the reduction of food consumption ($g = 0.560$, CI: 0.279–0.841; $z = 3.908$, $p < 0.0001$). No study has investigated the effect of neuromodulation on consumption in individuals with alcohol or illicit drug addiction.

Comparing different stimulation techniques

The comparison of the two stimulation techniques (rTMS vs. tDCS) showed no significant differences in their overall




Fig. 8. The effect of multi-session neuromodulation on consumption.

effectiveness for down-regulating craving ($Q = 0.307$, $p = 0.579$); with a significant effect for tDCS ($g = 0.490$, CI: 0.347–0.633; $z = 6.710$, $p < 0.0001$) and rTMS interventions ($g = 0.411$, CI: 0.170–0.651; $z = 3.350$, $p = 0.001$). When separately considering the effect on craving within single-session ($Q = 0.398$, $p = 0.528$) or multi-session interventions ($Q = 0.054$, $p = 0.816$), we also found no significant differences in the effectiveness between the two stimulation techniques. There were not enough studies to assess the session effect for consumption.

Comparing different stimulated hemispheres

The comparison of the overall effect size depending on the side of stimulation (right versus left hemisphere) revealed no significant differences for an effect on craving ($Z = 0.028$, $p = 0.9778$), with similar effects for each hemisphere separately (left: $g = 0.438$, CI: 0.254–0.623; $z = 4.651$, $p < 0.0001$; right: $g = 0.464$, CI: 0.300–0.628; $z = 5.545$, $p < 0.0001$). This was replicated when considering only single-session ($Z = 0.01$, $p = 0.992$) or multi-session interventions ($Z = 0.088$, $p = 0.930$). Similarly, the side of stimulation did not influence the overall effectiveness of the intervention in reducing consumption levels ($Z = 0.045$, $p = 0.964$), with similar effects per targeted hemisphere (left: $g = 0.602$, CI: 0.305–0.899; $z = 3.966$, $p < 0.0001$; right: $g = 0.648$, CI: 0.220–1.076; $z = 2.968$, $p = 0.003$).

Influence of study design

Finally, when investigating the influence of the chosen study design, we found a significant craving reduction both in studies with a between-subject design ($g = 0.541$, CI: 0.332–0.750; $z = 5.078$, $p < 0.0001$) and a within-subject design ($g = 0.399$, CI: 0.237–0.561; $z = 4.828$, $p < 0.0001$); there was no significant difference in the effect size between the two study designs ($Q = 1.114$, $p = 0.291$). Consistent with this pattern, we found a significant reduction of consumption levels in studies using a between-subject design ($g = 0.690$, CI: 0.346–1.034; $z = 3.930$, $p < 0.0001$) or a within-subject design ($g = 0.645$, CI: 0.261–1.029; $z = 3.289$, $p = 0.001$); and again no difference in the effect size between the two designs ($Q = 0.029$, $p = 0.865$).

Discussion

The current meta-analysis of 48 studies revealed a significant overall effect of neuromodulation interventions targeted at the dlPFC (over sham stimulation), with small to medium effect sizes for the reduction of craving and small to large effect sizes for the reduction of consumption levels across different addicted populations. More importantly, the current results showed for the first

time that multi-session protocols are more effective than single-session protocols. We found significantly larger effects of multi-session interventions both on craving reduction and for ability to refrain from consumption, with a linear dose-response effect such that an increased number of sessions or administered pulses had a larger effect on the craving level.

These results are in agreement with the general pattern of prolonged neuromodulation: compared to single pulse TMS, repeated pulses induce more prolonged effects in the brain, an effect which scales with the number of pulses applied [37]. Moreover, multi-session excitatory rTMS with its multifold total number of pulses has more prolonged effects than single-session rTMS does, generating a cumulative long-term potentiation (LTP) of synaptic connections [37]. Similarly, the length of stimulation in tDCS neuromodulation is crucial to achieving sustained effects [39]. The current meta-analytical results are also consistent with previous individual studies showing a larger effect of multi-session as compared to single-session interventions [45–47,65,66], which reported a linear dose-response effect of neuromodulation [46]. While we would expect a saturation of this dose-response effect after a certain number of sessions, the current results indicate that this threshold has not been reached in the reviewed studies.

The current results hence imply that multi-session excitatory stimulation protocols targeted at the dlPFC may have a cumulative dose effect on craving and consumption, potentially improving inhibitory control capacity underlying successful self-regulation of craving and consumption through normalizing hypo-active dlPFC functions during attempted self-regulation. As the current results suggest that neuromodulation interventions are equally effective in individuals with drug addiction and the ones with eating addiction, this meta-analysis also provides evidence for the hypothesized similarity in the underlying neuro-behavioral mechanisms, though differences across these populations surely exist [16]. It is interesting to note that we replicated all findings within a restricted sample of potentially more impaired individuals (excluding individuals without a clinical diagnosis), demonstrating the general clinical benefit of the intervention across very different populations.

The current results were also in agreement with previous meta-analyses, which showed small-sized effects on craving reduction across different addicted populations [41], a medium-sized effect for nicotine [44] and food [42]. However, in contrast to Maiti's previous meta-analysis that found no significant effect on alcohol craving [44], the current analysis also showed a small effect in this population. This discrepancy may be due to the limited number of studies (6 studies) in Maiti's meta-analysis versus a relatively larger number of studies in our meta-analysis (12 studies). The current results also showed for the first time that there was a significant

effect (medium-sized) for a craving reduction in illicit drug users (e.g., cannabis, cocaine, heroin, methamphetamine) by brain stimulation, hence suggesting consistent effects across all four included populations (alcohol vs. nicotine vs. illicit drugs vs. eating addictions).

Moreover, regarding the effect of neuromodulation on consumption, the current analysis (covering 12 studies) is consistent with a previously updated meta-analysis (covering 8 studies) focusing on individuals with excessive eating, which found an effect of brain stimulation on food consumption [43]. Furthermore, the current results demonstrated for the first time that non-invasive brain stimulation targeted at dlPFC can lead to a large-sized effect in the consumption reduction of cigarette smoking. When comparing smokers to excessive eaters, we found a marginally significant difference in the effect sizes for reduction of consumption, with a large effect size for smokers and a medium effect size for eating addiction. However, this difference between the reduction in cigarette and food consumption may be in part due to differences in the used measurement instruments, as the reduction in cigarette consumption was mainly assessed by self-report whereas the reduction in food taking was generally assessed with the real consumption (e.g., the amounts of calories consumed) after the intervention. There was no difference in effect sizes between populations when craving was used as an outcome measure.

The current results, also converging with a previous meta-analysis of 17 studies [41], showed that the stimulation technique *per se* did not significantly affect the reduction of craving across different populations. This pattern was different from a previous meta-analysis (covering 12 studies, including 8 tDCS) [42,43] which showed a stimulation effect on food cravings that was statistically significant for rTMS but not tDCS. When we restricted our analysis to studies on food craving only (covering 16 studies, including 11 for tDCS), we showed a significant brain stimulation effect for both rTMS ($g = 0.287$, CI: 0.014–0.559; $z = 2.061$, $p = 0.039$) and tDCS ($g = 0.479$, CI: 0.266–0.692; $z = 4.400$, $p < 0.001$).

In the current meta-analysis, we examined the potential effect of lateralized neuromodulation targeted at a specific hemisphere (left versus right) but found no effect for this factor, consistent with a previous meta-analysis [41], but inconsistent with a second meta-analysis, which only investigated the effect on food craving [43]. When restricting our analysis to include only studies on food craving we did not find a difference between the targeted site ($Z = 0.170$, $p = 0.865$). However, it would be premature to conclude that the mechanism of intervention may not be lateralized, as there are strong anatomical connections between the left and right dlPFC, likely leading to the presence of neuromodulation in both hemispheres. Finally, we examined the potential effect of study design (within-versus between-participant) for the first time, and again found no effect of this variable, confirming the independence of our results from these operational parameters.

Despite its strengths, the current meta-analysis has several limitations. First, the currently available literature does not allow for a systematic investigation of long-term outcomes of brain stimulation, as studies with a follow-up visit were scarce. The few studies that did include long-term outcome measures had somewhat inconsistent findings: some studies observed a long-term reduction of craving 10 days [72], 25 days [73] or one month [74] after the last brain stimulation of dlPFC or a long-term reduction of consumption nine days [65], 14–21 days [75] or six months [76] after the last stimulation; other studies observed no such long-term effects on craving either 1–3 days [77] or six months [78] after the last treatment. Second, due to the large heterogeneity in the used stimulation protocols, we were not able to evaluate the potential effects of many variables involved in the protocols. For example, we

could not explore whether the frequency (10 versus 20 Hz) used in rTMS protocols influence the modulation effects, as only five studies used 20 Hz protocols [77,79–82]. Similarly, we could not evaluate the influence of the electric current intensity used in tDCS studies (1 vs 2 mA), as only four studies adopted an 1 mA protocol [47,83–85]. In the current meta-analysis we further excluded studies using (inhibitory) low-frequency rTMS (≤ 1 Hz). In contrast to excitatory high frequency (≥ 5 Hz) rTMS intervention, low frequency rTMS studies targeted at dlPFC have been shown to increase craving levels in methamphetamine users [86], however, other studies have found opposite results in methamphetamine users [87] and pathological gamblers [88]. These discrepancies need to be further explored, as only a limited number of studies is currently available. Third, we only focused on excitatory brain stimulation of the dlPFC, studies that targeted at brain regions other than dlPFC (e.g., medial PFC [89]; cathodal frontal-parietal-temporal [90]) were not evaluated due to the limited numbers of relevant studies. Initial pilot studies employing inhibitory forms of TMS (cTBS = continuous bursting frequency TMS) targeting ventral medial prefrontal cortex (vmPFC) demonstrated the feasibility of this approach for attenuating reactivity of the reward network and reducing craving [91,92]. Overall, the optimization of stimulation protocols and definition of neural targets certainly remain an important goal for future research.

Conclusions

The current meta-analysis found significant effects of neuromodulation approaches targeted at the dlPFC for reducing craving and consumption in both eating and drug addiction. Importantly, multi-session interventions and longer sessions had larger effects than single-session interventions, with a linear dose-response effect. Overall, the current findings support the efficacy of neuromodulation approaches as a clinical treatment for individuals with drug or eating addiction.

Conflict-of-interest statement

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Acknowledgements

This work was supported by the National Basic Research Program of China (973 Program: 2015CB856400) and National Natural Science Foundation of China (31630034).

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.brs.2018.12.975>.

References

- [1] Center for Behavioral Health Statistics and Quality. 2016 national Survey on drug use and health: detailed tables. Rockville, MD: Substance Abuse and Mental Health Services Administration; 2017.
- [2] National Center for Health Statistics. Health, United States, 2016: with chartbook on long-term trends in health. Hyattsville, Maryland. 2017.
- [3] Hudson JI, Hiripi E, Pope HG, Kessler RC. The prevalence and correlates of eating disorders in the National Comorbidity Survey Replication. *Biol Psychiatry* 2007;61(3):348–58.
- [4] National Institute on Drug Abuse. Principles of drug addiction treatment. 3th ed. 2012. https://www.drugabuse.gov/sites/default/files/podat_1.pdf. [Accessed 20 August 2018].
- [5] Kass AE, Kolko RP, Wilfley DE. Psychological treatments for eating disorders. *Curr Opin Psychiatr* 2013;26(6):549.

- [6] Association AP. Diagnostic and statistical manual of mental disorders. fifth ed. Washington, DC: American Psychiatric Association; 2013.
- [7] Tiffany ST, Wray JM. The clinical significance of drug craving. *Ann Ny Acad Sci* 2012;1248(1):1–17.
- [8] Wang G-J, Volkow ND, Thanos PK, Fowler JS. Similarity between obesity and drug addiction as assessed by neurofunctional imaging: a concept review. *J Addict Dis* 2004;23(3):39–53.
- [9] Potenza MN, Grilo CM. How relevant is food craving to obesity and its treatment? *Front Psychiatr* 2014;5:164.
- [10] Chao A, Grilo CM, White MA, Sinha R. Food cravings, food intake, and weight status in a community-based sample. *Eat Behav* 2014;15(3):478–82.
- [11] Goldstein RZ, Volkow ND. Dysfunction of the prefrontal cortex in addiction: neuroimaging findings and clinical implications. *Nat Rev Neurosci* 2011;12(11):652.
- [12] Kaye WH, Wierenga CE, Bailer UF, Simmons AN, Wagner A, Bischoff-Grethe A. Does a shared neurobiology for foods and drugs of abuse contribute to extremes of food ingestion in anorexia and bulimia nervosa? *Biol Psychiatry* 2013;73(9):836–42.
- [13] Volkow ND, Wang G-J, Tomasi D, Baler RD. The addictive dimensionality of obesity. *Biol Psychiatry* 2013;73(9):811–8.
- [14] Gearhardt AN, White MA, Potenza MN. Binge eating disorder and food addiction. *Curr Drug Abuse Rev* 2011;4(3):201–7.
- [15] Davis C, Carter JC. Compulsive overeating as an addiction disorder. A review of theory and evidence. *Appetite* 2009;53(1):1–8.
- [16] Hebebrand J, Albayrak O, Adan R, Antel J, Dieguez C, de Jong J, et al. "Eating addiction", rather than "food addiction", better captures addictive-like eating behavior. *Neurosci Biobehav Rev* 2014;47:295–306.
- [17] Everitt BJ, Robbins TW. Drug addiction: updating actions to habits to compulsions ten years on. *Annu Rev Psychol* 2016;67:23–50.
- [18] Koob GF, Volkow ND. Neurobiology of addiction: a neurocircuitry analysis. *Lancet Psychiatry* 2016;3(8):760–73.
- [19] Koob GF, Volkow ND. Neurocircuitry of addiction. *Neuropsychopharmacology* 2010;35(1):217.
- [20] Robinson TE, Berridge KC. The neural basis of drug craving: an incentive-sensitization theory of addiction. *Brain Res Rev* 1993;18(3):247–91.
- [21] Belin D, Belin-Rauscent A, Murray JE, Everitt BJ. Addiction: failure of control over maladaptive incentive habits. *Curr Opin Neurobiol* 2013;23(4):564–72.
- [22] Goldstein RZ, Volkow ND. Drug addiction and its underlying neurobiological basis: neuroimaging evidence for the involvement of the frontal cortex. *Am J Psychiatry* 2002;159(10):1642–52.
- [23] Goldman M, Szucs-Reed RP, Jagannathan K, Ehrman RN, Wang Z, Li Y, et al. Reward-related brain response and craving correlates of marijuana cue exposure: a preliminary study in treatment-seeking marijuana-dependent subjects. *J Addiction Med* 2013;7(1):8.
- [24] Kühn S, Gallinat J. Common biology of craving across legal and illegal drugs—a quantitative meta-analysis of cue-reactivity brain response. *Eur J Neurosci* 2011;33(7):1318–26.
- [25] Jentsch JD, Taylor JR. Impulsivity resulting from frontostriatal dysfunction in drug abuse: implications for the control of behavior by reward-related stimuli. *Psychopharmacology* 1999;146(4):373–90.
- [26] Zilverstand A, Huang AS, Alia-Klein N, Goldstein RZ. Neuroimaging impaired response inhibition and salience attribution in human drug addiction: a systematic review. *Neuron* 2018;98(5):886–903.
- [27] Mansouri FA, Tanaka K, Buckley MJ. Conflict-induced behavioural adjustment: a clue to the executive functions of the prefrontal cortex. *Nat Rev Neurosci* 2009;10(2):141.
- [28] Koehlin E, Hyafil A. Anterior prefrontal function and the limits of human decision-making. *Science* 2007;318(5850):594–8.
- [29] Badre D. Cognitive control, hierarchy, and the rostral-caudal organization of the frontal lobes. *Trends Cognit Sci* 2008;12(5):193–200.
- [30] LuijtenBadre D, Nee DE. Frontal cortex and the hierarchical control of behavior. *Trends Cognit Sci* 2018;22(2):170–88.
- [31] Luijten M, Machielsen MW, Veltman DJ, Hester R, de Haan L, Franken IH. Systematic review of ERP and fMRI studies investigating inhibitory control and error processing in people with substance dependence and behavioural addictions. *J Psychiatry Neurosci* 2014;39(3):149–69.
- [32] Zilverstand A, Parvaz MA, Moeller SJ, Goldstein RZ. Cognitive interventions for addiction medicine: understanding the underlying neurobiological mechanisms. In: *Progress in brain research*. Elsevier; 2016. p. 285–304.
- [33] Zilverstand A, Parvaz MA, Goldstein RZ. Neuroimaging cognitive reappraisal in clinical populations to define neural targets for enhancing emotion regulation. A systematic review. *Neuroimage* 2017;151:105–16.
- [34] Lowe CJ, Hall PA, Staines WR. The effects of continuous theta burst stimulation to the left dorsolateral prefrontal cortex on executive function, food cravings, and snack food consumption. *Psychosom Med* 2014;76(7):503–11.
- [35] Lowe CJ, Staines WR, Manocchio F, Hall PA. The neurocognitive mechanisms underlying food cravings and snack food consumption. A combined continuous theta burst stimulation (ctBS) and EEG study. *Neuroimage* 2018;177:45–58.
- [36] Spagnolo PA, Goldman D. Neuromodulation interventions for addictive disorders: challenges, promise, and roadmap for future research. *Brain* 2017;140(5):1183–203.
- [37] Ridding MC, Rothwell JC. Is there a future for therapeutic use of transcranial magnetic stimulation? *Nat Rev Neurosci* 2007;8(7):559.
- [38] Peterchev AV, Wagner TA, Miranda PC, Nitsche MA, Paulus W, Lisanby SH, et al. Fundamentals of transcranial electric and magnetic stimulation dose: definition, selection, and reporting practices. *Brain Stimul* 2012;5(4):435–53.
- [39] Nitsche MA, Paulus W. Sustained excitability elevations induced by transcranial DC motor cortex stimulation in humans. *Neurology* 2001;57(10):1899–901.
- [40] Brunoni AR, Ferrucci R, Fregni F, Boggio PS, Priori A. Transcranial direct current stimulation for the treatment of major depressive disorder: a summary of preclinical, clinical and translational findings. *Prog Neuro-Psychopharmacol Biol Psychiatry* 2012;39(1):9–16.
- [41] Jansen JM, Daams JG, Koeter MW, Veltman DJ, van den Brink W, Goudriaan AE. Effects of non-invasive neurostimulation on craving: a meta-analysis. *Neurosci Biobehav Rev* 2013;37(10 Pt 2):2472–80.
- [42] Lowe CJ, Vincent C, Hall PA. Effects of noninvasive brain stimulation on food cravings and consumption: a meta-analytic review. *Psychosom Med* 2017;79(1):2–13.
- [43] Hall PA, Lowe C, Vincent C. Brain stimulation effects on food cravings and consumption: an update on Lowe et al.(2017) and a response to Generoso et al.(2017). *Psychosom Med* 2017;79(7):839–42.
- [44] Maiti R, Mishra BR, Hota D. Effect of high-frequency transcranial magnetic stimulation on craving in substance use disorder: a meta-analysis. *J Neuropsychiatry Clin Neurosci* 2016;29(2):160–71.
- [45] Su H, Zhong N, Gan H, Wang J, Han H, Chen T, et al. High frequency repetitive transcranial magnetic stimulation of the left dorsolateral prefrontal cortex for methamphetamine use disorders: a randomised clinical trial. *Drug Alcohol Depend* 2017;175:84–91.
- [46] Boggio PS, Liguori P, Sultani N, Rezende L, Fecteau S, Fregni F. Cumulative priming effects of cortical stimulation on smoking cue-induced craving. *Neurosci Lett* 2009;463(1):82–6.
- [47] Jauch-Chara K, Kistenmacher A, Herzog N, Schwarz M, Schweiger U, Oltmanns KM. Repetitive electric brain stimulation reduces food intake in humans. *Am J Clin Nutr* 2014;100(4):1003–9.
- [48] Grall-Bronnec M, Sauvaget A. The use of repetitive transcranial magnetic stimulation for modulating craving and addictive behaviours: a critical literature review of efficacy, technical and methodological considerations. *Neurosci Biobehav Rev* 2014;47:592–613.
- [49] Sauvaget A, Trojak B, Bulteau S, Jiménez-Murcia S, Fernández-Aranda F, Wolz I, et al. Transcranial direct current stimulation (tDCS) in behavioral and food addiction: a systematic review of efficacy, technical, and methodological issues. *Front Neurosci* 2015;9:349.
- [50] Trojak B, Sauvaget A, Fecteau S, Lalanne L, Chauvet-Gelinier J-C, Koch S, et al. Outcome of non-invasive brain stimulation in substance use disorders: a review of randomized sham-controlled clinical trials. *J Neuropsychiatry Clin Neurosci* 2017;29(2):105–18.
- [51] Lupi M, Martinotti G, Santacroce R, Cinosi E, Carlucci M, Marini S, et al. Transcranial direct current stimulation in substance use disorders: a systematic review of scientific literature. *J ECT* 2017;33(3):203–9.
- [52] Dunlop K, Hanlon CA, Downar J. Noninvasive brain stimulation treatments for addiction and major depression. *Ann Ny Acad Sci* 2017;1394(1):31–54.
- [53] Hall PA, Vincent CM, Burhan AM. Non-invasive brain stimulation for food cravings, consumption, and disorders of eating: a review of methods, findings and controversies. *Appetite* 2018;124:78–88.
- [54] Claudino A, Van den Eynde F, Stahl D, Dew T, Andiappan M, Kalthoff J, et al. Repetitive transcranial magnetic stimulation reduces cortisol concentrations in bulimic disorders. *Psychol Med* 2011;41(6):1329.
- [55] Shahbabaie A, Golesorkhi M, Zamanian B, Ebrahimipoor M, Keshvari F, Nejati V, et al. State dependent effect of transcranial direct current stimulation (tDCS) on methamphetamine craving. *Int J Neuropsychopharmacol* 2014;17(10):1591–8.
- [56] Boggio PS, Sultani N, Fecteau S, Merabet L, Mecca T, Pascual-Leone A, et al. Prefrontal cortex modulation using transcranial DC stimulation reduces alcohol craving: a double-blind, sham-controlled study. *Drug Alcohol Depend* 2008;92(1–3):55–60.
- [57] Boggio PS, Zaghi S, Villani AB, Fecteau S, Pascual-Leone A, Fregni F. Modulation of risk-taking in marijuana users by transcranial direct current stimulation (tDCS) of the dorsolateral prefrontal cortex (DLPFC). *Drug Alcohol Depend* 2010;112(3):220–5.
- [58] Fregni F, Orsati F, Pedrosa W, Fecteau S, Tome FA, Nitsche MA, et al. Transcranial direct current stimulation of the prefrontal cortex modulates the desire for specific foods. *Appetite* 2008;51(1):34–41.
- [59] Fregni F, Liguori P, Fecteau S, Nitsche MA, Pascual-Leone A, Boggio PS. Cortical stimulation of the prefrontal cortex with transcranial direct current stimulation reduces cue-provoked smoking craving: a randomized, sham-controlled study. *J Clin Psychiatr* 2008;69(1):32–40.
- [60] Kekic M, McClelland J, Bartholdy S, Boysen E, Musiat P, Dalton B, et al. Single-session transcranial direct current stimulation temporarily improves symptoms, mood, and self-regulatory control in bulimia nervosa: a randomised controlled trial. *PLoS One* 2017;12(1):e0167606.
- [61] Mitchell M, Muftakhidinov B, Winchen T, Jędrzejewski-Szmek Z. Engage digitizer software. Webpage: <http://markummitcheh.github.io/engage-digitizer>, Accessed 2017;11.
- [62] Higgins J, Green S. Handbook for systematic reviews of interventions version 5.1. 0 [updated March 2011]. The Cochrane Collaboration; 2011.

- [63] Borenstein M, Hedges LV, Higgins JP, Rothstein HR. Introduction to meta-analysis. 2009.
- [64] Borenstein M, Hedges LV, Higgins J, Rothstein HR. A basic introduction to fixed-effect and random-effects models for meta-analysis. *Res Synth Methods* 2010;1(2):97–111.
- [65] Fecteau S, Agosta S, Hone-Blanchet A, Fregni F, Boggio P, Ciraulo D, et al. Modulation of smoking and decision-making behaviors with transcranial direct current stimulation in tobacco smokers: a preliminary study. *Drug Alcohol Depend* 2014;140:78–84.
- [66] Shen Y, Cao X, Tan T, Shan C, Wang Y, Pan J, et al. 10-Hz repetitive transcranial magnetic stimulation of the left dorsolateral prefrontal cortex reduces heroin cue craving in long-term addicts. *Biol Psychiatry* 2016;80(3):E13–4.
- [67] Lipsey MW, Wilson DB. Practical meta-analysis. Sage Publications, Inc; 2001.
- [68] Egger M, Smith GD, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;315(7109):629–34.
- [69] Van den Eynde F, Claudino AM, Mogg A, Horrell L, Stahl D, Ribeiro W, et al. Repetitive transcranial magnetic stimulation reduces cue-induced food craving in bulimic disorders. *Biol Psychiatry* 2010;67(8):793–5.
- [70] Goldman RL, Borckardt JJ, Frohman HA, O'Neil PM, Madan A, Campbell LK, et al. Prefrontal cortex transcranial direct current stimulation (tDCS) temporarily reduces food cravings and increases the self-reported ability to resist food in adults with frequent food craving. *Appetite* 2011;56(3):741–6.
- [71] Sahlem GL, Baker NL, George MS, Malcolm RJ, McRae-Clark AL. Repetitive transcranial magnetic stimulation (rTMS) administration to heavy cannabis users. *Am J Drug Alcohol Abuse* 2017:1–9.
- [72] Bravo GL, Poje AB, Perissinotti I, Marcondes BF, Villamar MF, Manzardo AM, et al. Transcranial direct current stimulation reduces food-craving and measures of hyperphagia behavior in participants with Prader-Willi syndrome. *Am J Med Genet B Neuropsychiatr Genet* 2016;171B(2):266–75.
- [73] Ljubisavljevic M, Maxood K, Bjekic J, Oommen J, Nagelkerke N. Long-term effects of repeated prefrontal cortex transcranial direct current stimulation (tDCS) on food craving in normal and overweight young adults. *Brain Stimul* 2016;9(6):826–33.
- [74] Mishra BR, Nizamie SH, Das B, Praharaj SK. Efficacy of repetitive transcranial magnetic stimulation in alcohol dependence: a sham-controlled study. *Addiction* 2010;105(1):49–55.
- [75] Huang W, Shen F, Zhang J, Xing B. Effect of repetitive transcranial magnetic stimulation on cigarette smoking in patients with schizophrenia. *Shanghai Arch Psychiatry* 2016;28(6):309–17.
- [76] Dinur-Klein L, Dannon P, Hadar A, Rosenberg O, Roth Y, Kotler M, et al. Smoking cessation induced by deep repetitive transcranial magnetic stimulation of the prefrontal and insular cortices: a prospective, randomized controlled trial. *Biol Psychiatry* 2014;76(9):742–9.
- [77] Herremans S, Baeken C, Vanderbruggen N, Vanderhasselt M-A, Zeeuws D, Santermans L, et al. No influence of one right-sided prefrontal HF-rTMS session on alcohol craving in recently detoxified alcohol-dependent patients: results of a naturalistic study. *Drug Alcohol Depend* 2012;120(1):209–13.
- [78] Amiaz R, Levy D, Vainiger D, Grunhaus L, Zangen A. Repeated high-frequency transcranial magnetic stimulation over the dorsolateral prefrontal cortex reduces cigarette craving and consumption. *Addiction* 2009;104(4):653–60.
- [79] Wing VC, Bacher I, Wu BS, Daskalakis ZJ, George TP. High frequency repetitive transcranial magnetic stimulation reduces tobacco craving in schizophrenia. *Schizophr Res* 2012;139(1):264–6.
- [80] Höppner J, Broese T, Wendler L, Berger C, Thome J. Repetitive transcranial magnetic stimulation (rTMS) for treatment of alcohol dependence. *World J Biol Psychiatr* 2011;12(sup 1):57–62.
- [81] Herremans S, Vanderhasselt M-A, De Raedt R, Baeken C. Reduced intra-individual reaction time variability during a Go–NoGo task in detoxified alcohol-dependent patients after one right-sided dorsolateral prefrontal HF-rTMS session. *Alcohol Alcohol* 2013;48(5):552–7.
- [82] Herremans SC, Van Schuerbeek P, De Raedt R, Matthys F, Buyl R, De Mey J, et al. The impact of accelerated right prefrontal high-frequency repetitive transcranial magnetic stimulation (rTMS) on cue-reactivity: an fMRI study on craving in recently detoxified alcohol-dependent patients. *PLoS One* 2015;10(8).
- [83] den Uyl TE, Gladwin TE, Wiers RW. Transcranial direct current stimulation, implicit alcohol associations and craving. *Biol Psychol* 2015;105:37–42.
- [84] Nakamura-Palacios EM, de Almeida Benevides MC, da Penha Zago-Gomes M, de Oliveira RWD, de Vasconcelos VF, de Castro LNP, et al. Auditory event-related potentials (P3) and cognitive changes induced by frontal direct current stimulation in alcoholics according to Lesch alcoholism typology. *Int J Neuropsychopharmacol* 2012;15(5):601–16.
- [85] Wietschorke K, Lippold J, Jacob C, Polak T, Herrmann MJ. Transcranial direct current stimulation of the prefrontal cortex reduces cue-reactivity in alcohol-dependent patients. *J Neural Transm* 2016;123(10):1173–8.
- [86] Li X, Malcolm RJ, Huebner K, Hanlon CA, Taylor JJ, Brady KT, et al. Low frequency repetitive transcranial magnetic stimulation of the left dorsolateral prefrontal cortex transiently increases cue-induced craving for methamphetamine: a preliminary study. *Drug Alcohol Depend* 2013;133(2):641–6.
- [87] Liu Q, Shen Y, Cao X, Li Y, Chen Y, Yang W, et al. Either at left or right, both high and low frequency rTMS of dorsolateral prefrontal cortex decreases cue induced craving for methamphetamine. *Am J Addict* 2017;26(8):776–9.
- [88] Sauvaget A, Bulteau S, Guilleux A, Leboucher J, Pichot A, Valriviere P, et al. Both active and sham low-frequency rTMS single sessions over the right DLPFC decrease cue-induced cravings among pathological gamblers seeking treatment: a randomized, double-blind, sham-controlled crossover trial. *J Behav Addict* 2018;7(1):126–36.
- [89] Ceccanti M, Inghilleri M, Attilia ML, Raccach R, Fiore M, Zangen A, et al. Deep TMS on alcoholics: effects on cortisolemia and dopamine pathway modulation. A pilot study. *Can J Physiol Pharmacol* 2015;93(4):283–90.
- [90] Wang Y, Shen Y, Cao X, Shan C, Pan J, He H, et al. Transcranial direct current stimulation of the frontal-parietal-temporal area attenuates cue-induced craving for heroin. *J Psychiatr Res* 2016;79:1–3.
- [91] Hanlon CA, Dowdle LT, Austelle CW, DeVries W, Mithoefer O, Badran BW, et al. What goes up, can come down: novel brain stimulation paradigms may attenuate craving and craving-related neural circuitry in substance dependent individuals. *Brain Res* 2015;1628:199–209.
- [92] Hanlon CA, Kearney-Ramos T, Dowdle LT, Hamilton S, DeVries W, Mithoefer O, et al. Developing repetitive transcranial magnetic stimulation (rTMS) as a treatment tool for cocaine use disorder: a series of six translational studies. *Curr Behav Neurosci Rep* 2017;4(4):341–52.